



Carbonylation

The development of efficient and environmentally benign methods for the synthesis of industrially relevant molecules is central to improving the green credentials of the pharmaceutical and fine chemicals industries. In this respect, catalysis is a key chemistry for achieving this. Of the broad arrays of catalytic reactions, the functionalisation of hydrocarbon feedstocks into more complex molecules using carbonylation is key area of research.

Carbonylation is formal addition of C=O unit into a molecule and the most atom efficient route to this is by using carbon monoxide (CO). Carbonylation is one of the core industrial technologies for the conversion of bulk materials into products of value for example, olefins can be converted by carbonylation to a variety of products such as alcohols, aldehydes and carboxylic acid derivatives. It has also been employed for the efficient synthesis of amides.

Despite its position as a core transformation industrially, carbonylations are rarely used in the synthesis of complex pharmaceutical molecules. Beller *et al.* have hypothesised that this may be due to a reluctance by the research communities to operate the high pressure equipment involved when running reaction involving gasses, though they note that a number of catalytic methods for carbonylations are carried out at ambient to low pressures. [1]

1. A. Brennführer, H. Neumann and M. Beller, **Palladium-Catalyzed Carbonylation Reactions of Aryl Halides and Related Compounds**, *Angew. Chem. Int. Ed.*, 2009, 48, 4114-4133.


This resource has been created as part of the IMI funded CHEM21 project (Chemical Manufacturing Methods for the 21st Century Pharmaceutical Industries). CHEM21 has received funding from the Innovative Medicines Initiative Joint Undertaking under grant agreement n°115360, resources of which are composed of financial contribution from the European Union's Seventh Framework Programme (FP7/2007-2013) and EFPIA companies' in kind contribution.

The educational material is licensed (unless otherwise specified) under Creative Commons license [CC BY-NC 4.0](#), which gives permission for it to be shared and adapted for non-commercial purposes as long as attribution is given. For full details please see our [legal statements](#).

The views expressed in regards to education and training materials represent the aspiration of the CHEM21 consortium, although may not always be the view of each individual organisation. Referencing of external sources does not imply formal endorsement by the CHEM21 consortium.

Case studies

Carbonylative α -arylations have seen a lot of interest over the years; however, initial methods were limited to the use of malonate derivatives as starting materials.[1][2][3] Catalytic systems for the intermolecular carbonylative α -arylation of ketones have been developed although these reactions were limited to the use of C-nucleophiles furnished with a carbonyl group.[4][5][6][7] Thus, extending carbonylative α -arylation towards nitriles as readily available coupling partners represents an attractive goal for further development. The resultant β -ketonitrile products from such methods are useful bifunctional intermediates for the synthesis of pharmaceutically relevant compounds.[8][9][10][11] As important optically active intermediates, the diversity of available β -ketonitriles has a significant impact on the range of structures that can be accessed.[12][13][14][15][16][17] As such, the conventional methods for the synthesis of β -ketonitriles through acylation of acetonitriles or carbonylative coupling of trimethylsilylacetonitrile (**Scheme 1**) are limited to the formation of β -ketones that are unsubstituted at the α -position.[18]




Scheme 1: β -ketonitrile formation from carbonylative coupling of trimethylsilylacetonitrile ([18])

Although recent approaches have been reported that enable access to α -substituted β -ketonitriles, these protocols do not allow for the generation of a quaternary α -carbon centre.[19][20] As a result, the development of protocol that allows access to β -ketonitriles from aryl halides, carbon monoxide and uses nitriles as a simple abundant feedstock represents an advantageous achievement. CHEM21 researchers have developed the first example of such a transformation using a commercially available catalyst system that allows for the selective formation of α -disubstituted β -ketonitriles under low CO pressures with the use of unactivated nitriles (**Scheme 2**).[21]

This resource has been created as part of the IMI funded CHEM21 project (Chemical Manufacturing Methods for the 21st Century Pharmaceutical Industries). CHEM21 has received funding from the Innovative Medicines Initiative Joint Undertaking under grant agreement n°115360, resources of which are composed of financial contribution from the European Union's Seventh Framework Programme (FP7/2007-2013) and EFPIA companies' in kind contribution.

The educational material is licensed (unless otherwise specified) under Creative Commons license [CC BY-NC 4.0](https://creativecommons.org/licenses/by-nc/4.0/), which gives permission for it to be shared and adapted for non-commercial purposes as long as attribution is given. For full details please see our [legal statements](#).

The views expressed in regards to education and training materials represent the aspiration of the CHEM21 consortium, although may not always be the view of each individual organisation. Referencing of external sources does not imply formal endorsement by the CHEM21 consortium.



Scheme 2: CHEM21 Approach and first example of the synthesis of α -disubstituted β -ketonitriles from unactivated nitriles([21])

The method is applicable to the synthesis of 24 α -disubstituted β -ketonitriles in good to excellent yields, which were achieved in a straightforward approach. The approach allows for the selective carbonylative α -arylation in an atom economical manner using CO gas.

1. E. -ichi Negishi, H. Makabe, I. Shimoyama, G. Wu and Y. Zhang, **Catalytic generation and trapping of acylmetals containing Ni and Cu with enolates**, *Tetrahedron*, 1998, **54**, 1095-1106.
2. E. -ichi Negishi, C. Copéret, T. Sugihara, I. Shimoyama, Y. Zhang, G. Wu and J. M. Tour, **Palladium-catalyzed carbonylative cyclization via trapping of acylpalladium derivatives with internal enolates. Its scope and factors affecting the C-to-O ratio**, *Tetrahedron*, 1994, **50**, 425-436.
3. Z. Zheng and H. Alper, **Palladium-Catalyzed Cyclocarbonylation-Decarboxylation of Diethyl(2-iodoaryl)malonates with Vinyl Ketones Affording Functionalized Enolic 2-Acyl-3,4-dihydronaphthalenones**, *Org. Lett.*, 2009, **11**, 3278-3281.
4. S. Korsager, D. U. Nielsen, R. H. Taaning and T. Skrydstrup, **Access to β -Keto Esters by Palladium-Catalyzed Carbonylative Coupling of Aryl Halides with Monoester Potassium Malonates**, *Angew. Chem.*, 2013, **125**, 9945-9948.
5. T. M. Gøgsig, R. H. Taaning, A. T. Lindhardt and T. Skrydstrup, **Palladium-Catalyzed Carbonylative α -Arylation for Accessing 1,3-Diketones**, *Angew. Chem. Int. Ed.*, 2012, **51**, 798-801.
6. D. U. Nielsen, C. Lescot, T. M. Gøgsig, A. T. Lindhardt and T. Skrydstrup, **Pd-Catalyzed Carbonylative α -Arylation of Aryl Bromides: Scope and Mechanistic Studies**, *Chem. Eur. J.*, 2013, **19**, 17926-17938.
7. J. Schranck, A. Tlili, H. Alsabeh Pamela G and Neumann, M. Stradiotto and M. Beller, **Palladium-catalysed carbonylative alpha-arylation of acetone and acetophenones to 1,3-diketones**, *Chemistry*, 2013, **19**, 12624-12628.
8. M. Ram Yadav, V. P. Pawar, S. M. Marvaniya, P. K. Halen, R. Giridhar and A. K. Mishra, **Site specific chemical delivery of NSAIDs to inflamed joints: Synthesis, biological activity and γ -imaging studies of quaternary ammonium salts of tropinol esters of some NSAIDs or their active metabolites**, *Biorg. Med. Chem.*,

This resource has been created as part of the IMI funded CHEM21 project (Chemical Manufacturing Methods for the 21st Century Pharmaceutical Industries). CHEM21 has received funding from the Innovative Medicines Initiative Joint Undertaking under grant agreement n°115360, resources of which are composed of financial contribution from the European Union's Seventh Framework Programme (FP7/2007-2013) and EFPIA companies' in kind contribution.

The educational material is licensed (unless otherwise specified) under Creative Commons license [CC BY-NC 4.0](https://creativecommons.org/licenses/by-nc/4.0/), which gives permission for it to be shared and adapted for non-commercial purposes as long as attribution is given. For full details please see our [legal statements](#).

The views expressed in regards to education and training materials represent the aspiration of the CHEM21 consortium, although may not always be the view of each individual organisation. Referencing of external sources does not imply formal endorsement by the CHEM21 consortium.

2008, **16**, 9443-9449.

9. I. Andreu, I. M. Morera, F. Bosca, L. Sanchez, P. Camps and M. A. Miranda, **Cholesterol-diaryl ketone stereoisomeric dyads as models for "clean" type I and type II photooxygenation mechanisms**, *Organic & Biomolecular Chemistry*, 2008, **6**, 860-867.
10. J. Norinder, K. Bogár, L. Kanupp and J. - E. Bäckvall, **An Enantioselective Route to α -Methyl Carboxylic Acids via Metal and Enzyme Catalysis**, *Org. Lett.*, 2007, **9**, 5095-5098.
11. A. Herschhorn, L. Lerman, M. Weitman, I. Oz Gleenberg, A. Nudelman and A. Hizi, **De Novo Parallel Design, Synthesis and Evaluation of Inhibitors against the Reverse Transcriptase of Human Immunodeficiency Virus Type-1 and Drug-Resistant Variants**, *J. Med. Chem.*, 2007, **50**, 2370-2384.
12. O. Soltani, M. A. Ariger, H. Vázquez-Villa and E. M. Carreira, **Transfer Hydrogenation in Water: Enantioselective, Catalytic Reduction of α -Cyano and α -Nitro Substituted Acetophenones**, *Org. Lett.*, 2010, **12**, 2893-2895.
13. D. Zhu, H. Ankati, C. Mukherjee, Y. Yang, E. R. Biehl and L. Hua, **Asymmetric Reduction of β -Ketonitriles with a Recombinant Carbonyl Reductase and Enzymatic Transformation to Optically Pure β -Hydroxy Carboxylic Acids**, *Org. Lett.*, 2007, **9**, 2561-2563.
14. H. Ankati, D. Zhu, Y. Yang, E. R. Biehl and L. Hua, **Asymmetric Synthesis of Both Antipodes of β -Hydroxy Nitriles and β -Hydroxy Carboxylic Acids via Enzymatic Reduction or Sequential Reduction/Hydrolysis**, *J. Org. Chem.*, 2009, **74**, 1658-1662.
15. P. Florey, A. J. Smallridge, A. Ten and M. A. Trehwella, **Chemo- and Stereoselective Reduction of an α -Cyanoketone by Bakers' Yeast at Low Temperature**, *Org. Lett.*, 1999, **1**, 1879-1880.
16. R. J. Hammond, B. W. Poston, I. Ghiviriga and B. D. Feske, **Biocatalytic synthesis towards both antipodes of 3-hydroxy-3-phenylpropanitrile a precursor to fluoxetine, atomoxetine and nisoxetine**, *Tetrahedron Lett.*, 2007, **48**, 1217-1219.
17. R. W. Nowill, T. J. Patel, D. L. Beasley, J. A. Alvarez, E. Jackson Iii, T. J. Hizer, I. Ghiviriga, S. C. Mateer and B. D. Feske, **Biocatalytic strategy toward asymmetric β -hydroxy nitriles and γ -amino alcohols**, *Tetrahedron Lett.*, 2011, **52**, 2440-2442.
18. A. Pa

This resource has been created as part of the IMI funded CHEM21 project (Chemical Manufacturing Methods for the 21st Century Pharmaceutical Industries). CHEM21 has received funding from the Innovative Medicines Initiative Joint Undertaking under grant agreement n°115360, resources of which are composed of financial contribution from the European Union's Seventh Framework Programme (FP7/2007-2013) and EFPIA companies' in kind contribution.

The educational material is licensed (unless otherwise specified) under Creative Commons license [CC BY-NC 4.0](#), which gives permission for it to be shared and adapted for non-commercial purposes as long as attribution is given. For full details please see our [legal statements](#).

The views expressed in regards to education and training materials represent the aspiration of the CHEM21 consortium, although may not always be the view of each individual organisation. Referencing of external sources does not imply formal endorsement by the CHEM21 consortium.